



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A NOTE ON THE CALCULATION OF EULER'S CONSTANT.

By GOLDIE HORTON, University of Texas.

Euler's constant, which plays an important rôle in the theory of gamma functions, is usually defined by the relation

$$\gamma = \lim_{n \rightarrow \infty} \sum_{m=1}^n \left[\frac{1}{m} - \log \left(1 + \frac{1}{m} \right) \right].$$

Its direct calculation from this definition is impracticable, and it is actually computed by means of the asymptotic expansion

$$\gamma = \lim_{n \rightarrow \infty} \sum_{m=1}^n \frac{1}{m} - \log n - \frac{1}{2n} + \frac{B_1}{2n^2} - \frac{B_2}{2n^4} + \dots, \quad (1)$$

where the B 's are the Bernoullian numbers, the error being less than the first term omitted.¹ The proof of (1) is a delicate matter, however, and it seems therefore worth while to call attention to the fact that γ can be calculated as readily as π or $\log n$ to three or four places of decimals, if one makes use of an idea employed in Cauchy's integral test. This method, which is applicable to all convergent series of positive monotone decreasing terms, depends upon the fact that from Cauchy's test it follows immediately that both

$$\sum_{m=1}^n u_m + \int_n^{\infty} u_m dm \quad \text{and} \quad \sum_{m=1}^n u_m + \int_{n+1}^{\infty} u_m dm$$

approximate $\sum_{m=1}^{\infty} u_m$ with an error less than $\int_n^{n+1} u_m dm$, and hence less than u_n .¹

Applying this to

$$\gamma = \sum_{m=1}^{\infty} \left[\frac{1}{m} - \log \left(1 + \frac{1}{m} \right) \right],$$

we see that

$$\sum_{m=1}^n \left[\frac{1}{m} - \log \left(1 + \frac{1}{m} \right) \right] + \int_n^{\infty} \left[\frac{1}{m} - \log \left(1 + \frac{1}{m} \right) \right] dm$$

or

$$\sum_{m=1}^n \left[\frac{1}{m} - \log \left(1 + \frac{1}{m} \right) \right] + \left[-1 - \log \left(\frac{n}{n+1} \right)^{n+1} \right]$$

approximates to Euler's constant with an error less than

$$\log \left(\frac{n+1}{n+2} \right)^{n+2} \left(\frac{n+1}{n} \right)^{n+1}$$

For $n = 5$, this gives $\gamma = .58249$ with an error less than .01487; for $n = 10$, $\gamma = .57948$ with an error less than .00429; for $n = 20$, $\gamma = .57779$ with an error less than .00115.

¹ See Wm. Shank's paper, "On the Calculation of the Numerical Value of Euler's Constant." *Proceedings of the Royal Society of London*, Vol. XV, p. 429.

¹ In these integrals the definition of the function u_m is extended so that it relates to the continuous variable m in such wise that u_m is a monotone decreasing function of the continuous variable m .